



Mo Turf

(0-0-0-0)
TL0057

FERTILIZERS

GUARANTEED ANALYSIS

Molybdenum (Mo)	23.7% w/v
Specific Gravity @ 20°C	1.398 kg/L
pH	7.0 to 8.0

20 L



APPLICATION RATES

GUIDING PRINCIPLES

Before using Mo Turf, seek appropriate agronomic advice via SLTEC's Balanced Agronomy services and dealer agronomist network.

Soil analysis combined with plant tissue (or SAP) analysis are recommended to assist in making an informed decision about which product best meets your crop, soil and general environmental conditions.

Fertilizer can burn and or damage crops and pasture. Because climatic and soil conditions, application methods, irrigation and management practices are beyond the control of Sustainable Liquid Technology Pty Ltd and cannot be foreseen, Sustainable Liquid Technology Pty Ltd accepts no responsibility whatsoever for any commercial damage, loss or other result following the use of this product, whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied or the rendering again of the services that are provided. The buyer accepts and uses this product subject to these conditions.

PRODUCT INFORMATION

PRODUCT STORAGE

Keep material in a shaded, cool, dry, well ventilated area, away from direct sunlight between 10 to 30°C. Store product in suitable containers made from material such as stainless steel, high density polyethylene and fibreglass.

Fluid fertilizers can be corrosive to material such as brass, copper, zinc or alloys of these metals. Clean equipment after use thoroughly with clean fresh water. Use SLTEC's Fert Tank Cleaner to rinse fertilizer tanks and transfer systems. Mo Turf is recommended for use within 12 months from dispatch date.

WARNING & SAFETY INFORMATION

WARNING: Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Wear rubber gloves and appropriate eye protection when handling the product. See the product SDS for more information, turfandlandscape.com.au

WARNING: Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of Molybdenum use with appropriate authorities.

Batch No. or Tank Number & Date:

Dispatch Date:f

PO Box 43 / 2055 Finlay Road
TONGALA VICTORIA 3621
P: 1800 768 224 F: 03 5859 1363
E: enquiries@sltec.com.au
ABN: 63 234 073 378 ACN: 113 670 269
www.sltec.com.au